

Case report

DYSPHAGIA AS AN EARLY PRESENTATION OF DIGEORGE'S SYNDROME – CASE REPORT

Snezana Palchevska¹, Beti Gurkova², Emilija Shukarova², Katarina Stavrikj², Jana Jovanovska², Natasa Aluloska²

¹ Clinical hospital *Adzibadem Sistina, Neonatology and Neonatal intensive care unit, , Skopje, Republic of North Macedonia*

² *University Clinic for Children's Diseases, Skopje; Ss Cyril and Methodius University, Faculty of Medicine, Skopje, Republic of North Macedonia*

Abstract

Citation: Palchevska S, Gurkova B, Shukarova E, Stavrikj K, Jovanovska J, Aluloska N. Dysphagia as an early presentation of DiGeorge's syndrome –case report.

doi.org/10.3889/aph.2021.5767

Key words: DiGeorge's syndrome, dysphagia, aberrant right subclavian artery, newborn.

***Correspondence:** Natasa Aluloska. University Clinic for children's diseases, Skopje, Republic of North Macedonia. E-mail: aluloska@gmail.com

Received: 7-Jan-2021; **Revised:** 20-Feb-2021; **Accepted:** 25-fev-2021; **Published:** 28-fev-2021

Copyright: © 2021. Snezana Palchevska, Beti Gurkova, Emilija Shukarova, Katarina Stavrikj, Jana Jovanovska, Natasa Aluloska. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Competing Interests: The author have declared that no competing interests

DiGeorge's syndrome is a 22q11.2 deletion leading to abnormal embryogenesis of pharyngeal arches and it is manifesting in a variety of clinical signs and symptoms. The spectrum of anomalies varies from minor facial dysmorphism and cleft palate to a broad spectrum of cardiovascular anomalies, thymic dysfunction and immune deficiencies, hypocalcemia due to hypoparathyroidism, growth and developmental delay and speech disturbances. Cardiovascular anomalies might include right sided aortic arch, aberrant vesicles and vascular ring. Here we present an atypical case of partial DiGeorge's syndrome with feeding and swallowing difficulties and laryngeal stridor in the neonatal period. Early presentation in this period is usually due to severe hypocalcemia and cardiac disease. Feeding difficulties in a preterm baby needed clinical assessment skills in order to establish the diagnosis and delineate it from feeding difficulties usually seen in preterm babies. Esophagogram (barium X Ray) showed antero-posterior oblique impression towards the right side, the latero- lateral view showed impression on the rare side, suspected to be esophageal sub stenosis due to vascular anomaly, aberrant right subclavian artery and suspected thymic hypoplasia. We report a 9-year follow up period by a team of subspecialists. The child had two surgeries due to aberrant vessel and velopharyngeal deficiency. Optimal management of patients with DiGeorge's syndrome requires a multidisciplinary team which should include a cardiologist, immunologist, geneticist, speech/language therapist, endocrinologist and other subspecialists depending on patient's phenotype.

Приказ на случај

ДИСФАГИЈА КАКО РАНА ПРЕЗЕНТАЦИЈА НА DIGEORGE СИНДРОМ - ПРИКАЗ НА СЛУЧАЈ

Снежана Палчевска¹, Бети Ѓуркова², Емилија Шукарова², Катарина Ставриќ², Јана Јовановска², Наташа Алулоска²

¹ *Клиничка болница Дџибадем Систина, Оддел за неонатологија и неонатална интензивна нега, Скопје, Република Северна Македонија*

² *Универзитетска Клиника за детски болести, Скопје; Универзитет Св. Кирил и Методиј, Скопје, Република Северна Македонија*

Изводок

Цитирање: Палчевска С, Ѓуркова Б, Шукарова Е, Ставриќ К, Јовановска Ј, Алулоска Н. Дисфагија како рана презентација на DiGeorge синдром - приказ на случај. Арх Ј Здравје 2021;13(1)

doi.org/10.3889/aph.2021.5767

Клучни зборови: Синдром на DiGeorge, дисфагија, aberrantна артерија субклавија, новородено.

***Кореспонденција:** Наташа Алулоска, Универзитетска клиника за детски болести, Скопје, Република Северна Македонија. E-mail: aluloska@gmail.com.

Примено: 7-јан-2021; **Ревидирано:** 20-фев-2021; **Прифатено:** 25-фев-2021; **Објавено:** 28-фев-2021

Печатарски права: ©2021 Снежана Палчевска, Бети Ѓуркова, Емилија Шукарова, Катарина Ставриќ, Јана Јовановска, Наташа Алулоска. Оваа статија е со отворен пристап дистрибуирана под условите на нелицензирана лиценца, која овозможува неограничена употреба, дистрибуција и репродукција на било кој медиум, доколку се цитираат оригиналниот(ите) автор(и) и изворот.

Конкурентски интереси: Авторот изјавува дека нема конкурентски интереси.

DiGeorge Синдромот е резултат на делеција на 22q11.2 која води до пореметена ембриогенеза на фарингеалните лази и се манифестира со различни знаци и симптоми. Спектарот на аномалии во овој синдром е варијабилен и може да се презентира само со минорна лицева дизморфија и расцеп на непцето но и со широк спектар на кардиоваскуларни аномалии, дисфункција на тимус и имун дефицит, хипокалцемија поради хипопаратиroidизам, пореметување во растот и развојот и проблеми во говорот. Кардиоваскуларните аномалии може да се од типот на десен аортен лак, aberrantни крвни садови и васкуларен ринг. Во овој приказ на случај прикажуваме атипичен случај на парцијален синдром на DiGeorge кој се презентираше со потешкотии во голтање и ларингеален стридор во неонаталниот период. Раната презентација на овој синдром во неонаталната возраст најчесто се дијагностицира со хипокалцемија и конгенитална кардиопатија. Потребна е добра клиничка проценка за да се разликува проблем при хранење на прематурно новородено од другите проблеми при голтање. Езофагограмот покажа коса импресија на антеро-постериорниот правец со ориентација кон десно суспектна за субстеноза на езофагусот на опишаното ниво од васкуларна етиологија, aberrantна десна артерија субклавија и суспектна хипоплазија на тимус. Пациентот беше следен во период од 9 години од тим на субспецијалисти. Во тој период имаше две хируршки интервенции (корекција на aberrantен крвен сад и велофарингеален дефицит). Пациентите со синдром на DiGeorge треба да бидат следени од мултидисциплинарен тим кој вклучува кардиолог, имунолог, генетичар, ендокринолог, логопед и други субспецијалисти според фенотипот на пациентот.

Introduction

DiGeorge's syndrome is a 22q11.2 deletion leading to abnormal embryogenesis of pharyngeal arches and it is manifesting in a variety of clinical signs and symptoms. The spectrum of anomalies varies from minor facial dysmorphism and cleft palate to a broad spectrum of cardiovascular anomalies, thymic dysfunction and immune deficiencies, hypocalcemia due to hypoparathyroidism, growth and developmental delay and speech disturbances. Cardiovascular anomalies include right sided aortic arch, aberrant vessels. When the aortic arch, or the associated vessels form a complete or incomplete ring around the trachea and the esophagus the condition is called vascular ring. It is rare congenital malformation that counts less than 1 % of all heart anomalies. The symptoms are recognised usually during the period of early infancy or later in life, because of feeding and respiratory difficulties.

We present an atypical case of partial DiGeorge's syndrome with feeding and swallowing difficulties and laryngeal stridor. Vascular ring should be suspected in all neonates with feeding and respiratory problems. Prompt diagnosis and treatment are of great importance for appropriate growth and development. In our case this condition is part of a syndrome, which should be carefully monitored by a multidisciplinary team.

Case report

One-month old infant was admitted to hospital due to feeding difficulties and vomiting. The condition

was worsening during feeding. The baby was born premature, 33 weeks gestational age, birth weight 1860 gr, Apgar score 7/8. Polyhydramnios was noticed in the last trimester. She was treated in the nursery for 3 weeks and was tube fed. On admission, neonatal infection was suspected, although the baby had normal values of inflammatory markers. The plain chest X-ray showed displaced mediastinal organs towards the right side, the upper right lobe had lower transparency suspected as consolidation. The tracheal aspirate was positive for *Klebsiella aerogenes* and antibiotic treatment was started. The baby had intermittent laryngeal stridor and feeding difficulties. Fiberlaryngoscopy was normal. Esophagogram (barium X Ray) showed antero-posterior oblique impression towards the right side, the latero-lateral view showed impression on the rare side, suspected to be esophageal stenosis due to vascular anomaly, aberrant right subclavian artery. Thymic hypoplasia was also suspected. The baby had normal calcium and parathormone levels. Echocardiography showed patent ductus arteriosus. Computer tomography with contrast showed vascular ring with left sided aortic arch with aberrant right subclavian artery and patent ductus arteriosus. Renal ultrasound was normal. FISH (Fluorescence in situ hybridization) was performed showing 22 q11.2 micro deletion (Figure 1).

At the age of 4 months the child had an operation to correct the vascular anomaly. Afterwards she had regular cardiologic assessments, showing normal cardiac structure and function. Following the procedure, swallowing status improved. The




Figure 1. FISH result:46, XX, del(22)(q11.2) FISH probe-Cytocell TUPLE1, green signal-control (22q13.3); red signal-locus-specific -(22q11.2)

growth remained beneath the third percentile for 9 years follow up. Neurodevelopmental assessment showed developmental delay with delayed speech development, learning delays and disabilities. ERA was normal. At the age of 3 years the child was diagnosed Velopharyngeal deficiency (VPD) by a speech and language pathologist. She was referred to Cleft lip and palate craniofacial clinic where VPD confirmed via nasoendoscopy. She was treated surgically with a posterior pharyngeal flap. She continued with speech therapy and was able to produce consonants for the first time and become intelligible to other people at the age of 5 years. Immunological assessment at age of 1 year showed normal function of the humoral immune system and neutrophil dysfunction. She was immunized by

inactivated vaccines regularly. Cellular immunity could not be accurately assessed, and immunization with live vaccine (MRP) was not performed. During the 9 year follow up she had mild upper respiratory infections and two episodes of bronchopneumonia that required hospital treatment at her early age.

Discussion

DiGeorge's syndrome features were first reported by Angelo DiGeorge in 1965 and 1968 although thymic aplasia was noted by Harrington in 1828 and the association with congenital hypoparathyroidism by Lobe in 1959^{1,2,3,4}. Additional anomalies were noted to be associated with the syndrome, especially anomalies of the great vessels, esophageal atresia, heart, ear, nose and mouth

defects⁵. For the first time the cause of the syndrome was suspected to be a deletion in chromosome 22q11 in 1981⁶. This is a 35 mb chromosomal region that contains more than 35 genes that affect the morphogenesis of the pharyngeal arches, heart and brain^{7,8,9,10}.

DiGeorge's syndrome is one of the most common genetic disorders. It occurs in 1:4000 live births but could be higher having in mind the clinical variability. More than 180 different defects have been associated with the syndrome¹¹. Typical presentation during infancy usually includes some of the following features: facial dysmorphia, heart defect, hypocalcemia, palatal anomalies and immunodeficiency¹². The standard method for diagnosis of DiGeorge's syndrome is the specific FISH test for chromosome 22q11 deletions. Due to the wide availability of these specific probes, clinicians can diagnose affected children, often presenting with cardiac abnormalities, immunodeficiency or hypocalcemic seizures¹³.

This syndrome is rarely diagnosed in the neonatal period. Early presentation in this period is usually due to severe hypocalcemia and cardiac disease¹⁴.

We present a rare case of early diagnosed patient in the neonatal period with feeding difficulties and swallowing problems. Our patient was a preterm baby, 33 weeks of gestation that stayed in the nursery for 3 weeks and was tube fed. She started having feeding difficulties when she was discharged home and was bottle fed. Feeding difficulties in a preterm baby needed clinical assessment skills in order to establish

the diagnosis and delineate it from feeding difficulties usually seen in preterm babies. The patient was admitted to our Clinic with aspiration-based infection as it has been previously reported^{15,16}.

Feeding difficulty is often a symptom of palatal dysfunction, cardiac anomalies and gastrointestinal dysmotility, as a result of the presence of a vascular ring that impairs the esophagus, and hypotonia can lead to feeding difficulty. In some patients, these difficulties likely result from craniofacial dysmorphology including cleft palate and other anomalies that often require surgical intervention^{7, 17,18}. Dysphagia can be seen in different developmental disorders including DiGeorge's syndrome¹⁹. As a symptom it may be present during infancy and may last with complications until the age of 4 years and sometimes onwards.

In our case, barium esophagogram showed esophageal sub stenosis due to vascular anomaly, aberrant right subclavian artery. And thymic hypoplasia or agenesis was suspected. The first report of symptomatic aberrant right subclavian artery was published in 1735, and in 1794 the term dysphagia lusoria was used to describe ARSA that has a retro-esophageal course^{20,21}.

Association of left sided aortic arch with aberrant right sided aortic arch and ipsilateral patent ductus arteriosus in a patient with DiGeorge's syndrome as seen in our patient was only described by Taliana et al²².

Conclusion

We present a rare case of DiGeorge's syndrome in a preterm infant with dysphagia due to aberrant right

subclavian artery. This was a case with a challenging diagnosis, in which classical manifestations of DiGeorge's syndrome was initially lacking. We report a 9-year follow up period by a team of subspecialists. The child had two surgeries due to aberrant vessel and velopharyngeal deficiency. Optimal management of patients with DiGeorge's syndrome requires a multidisciplinary team. The team should include a cardiologist, immunologist, geneticist, speech/language therapist, endocrinologist and other subspecialists depending on patient's phenotype, as reported in our case.

References

1. A comment on another paper, Cooper M., Peterson R., Good R. A new concept of the cellular basis of immunity. *The Journal of Pediatrics*.1965;67(5):907.
2. DiGeorge AM. Congenital absence of the thymus and its immunologic consequences: concurrence with congenital hypoparathyroidism. IV(1). White Plains, NY: March of Dimes-Birth Defects Foundation. 1968;116-21.
3. Harrington LH. Absence of the thymus gland. *Lond Med Gaz* 1929;3:314.
4. Lobdell DH. Congenital absence of the parathyroid glands. *Arch Pathol* 1959;67:412-18.
5. Lischner HW. DiGeorge syndrome(s). *Pediatr* 1972;81: 1042-4.
6. De la Chapelle A, Herva R, Koivisto M, Aula O. A deletion in chromosome 22 can cause DiGeorge syndrome. *Hum Genet* 1981;57:253-256.
7. Kobrynski LJ, Sullivan KE. Velocardiofacial syndrome, DiGeorge syndrome: the chromosome 22q11.2 deletion syndromes. *Lancet* 2007;370(9596):1443-1452.
8. Bassett AS, McDonald-McGinn DM, Devriendt K, et al; International 22q11.2 Deletion Syndrome Consortium. Practical guidelines for managing patients with 22q11.2 deletion syndrome. *J Pediatr* 2011;159(2):332-339.
9. Emanuel BS. Molecular mechanisms and diagnosis of chromosome 22q11.2 rearrangements. *Dev Disabil Res Rev* 2008;14(1):11-18.
10. Hurles M. How homologous recombination generates a mutable genome. *Hum Genomics* 2005;2(3):179-186.
11. Shprintzen RJ. Velo-cardio-facial syndrome. In: Cassidy SB, Allanson JE, editors. *Management of genetic syndromes*, 2nd ed. Hoboken: Wiley-Liss, 2005:615- 631.
12. McDonald-McGinn DM, Kirschner R, Goldmuntz E, Sullivan K, Eicher P, Gerdes M, et al. The Philadelphia story: the 22q11.2 deletion: report on 250 patients. *Genet Couns* 1999;10:11-24.
13. Yakut T, Kilic SS, Cil E, Yapici E, Egeli U. *Pediatr Surg Int* 2006; 22(4):380-3
14. Dogus Vuralli. Clinical approach to hypocalcemia in newborn period and infancy: Who should be treated? *International Journal of Pediatrics* 2019, Article ID 4318075
15. Hopkin RJ, Schorry EK, Bofinger M, Saal HM. Increased need for medical interventions in infants with velocardiofacial (deletion 22q11) syndrome. *J Pediatr* 2000; 137: 247-249.
16. Trinick R, Johnston N, Dalzell AM,

- McNamara P S. Reflux aspiration in children with neurodisability – a significant problem, but can we measure it? *J Pediatr Surg* 2012; 47, 291–298.
17. Oskarsdo'ttir S, Vujic M, Fasth A. Incidence and prevalence of the 22q11 deletion syndrome: a population-based study in Western Sweden. *Arch Dis Child* 2004;89:148–151.
 18. Botto LD, May K, Fernhoff PM, et al. A population-based study of the 22q11.2 deletion: phenotype, incidence, and contribution to major birth defects in the population. *Pediatrics* 2003;112:101–107.
 19. Eicher P.S. et al. Dysphagia in children with a 22q11.2 deletion: unusual pattern found on modified barium swallow. *J Pediatr* 2000; 137 (2):158–164.
 20. Williams GD, Schmeckebier M, Edmonds H W, Grand E G. Variations in the arrangement of the branches arising from the aortic arch in the American whites and negroes. *The Anatomical Record* 1932; 54: 247–251.
 21. Bayford D. An account of a singular case of obstructed deglutition. *Memoirs of the Medical Society of London* 1794; 2: 275–286.
 22. Taliana N, Gatt A, Borg A, Grech V. The rarest aortic arch anomaly a case report of asymptomatic isolation of the subclavian artery. *Images Paediatr Cardiol* 2017;19(2):9–12.