

CASE REPORT

MORNING GLORY SYNDROME

Igor Isjanovski¹, Bisera Velkovska¹¹ University Clinic for Eye Diseases; Ss Cyril and Methodius University in Skopje, Faculty of Medicine, Republic of North Macedonia**Abstract**

Morning Glory Syndrome (MGS) is an uncommon congenital anomaly of the eye nerve (optic nerve) that resembles a flower known as morning glory, impairs vision, and may be associated with both ocular and non-ocular abnormalities. It has a characteristic fundoscopic appearance consisting of a large funnel-shaped cavity on the optical disc. We register an unusual congenital anomaly of the optic disc in a three-year-old female child.

Citation: Isjanovski I, Velkovska B. Morning Glory Syndrome- A case report. Arch Pub Health 2022;14(1) 157-161.

doi.org/10.3889/aph.2022.6048

Key words: Morning Glory Syndrome, optical disc

***Correspondence:** Igor Isjanovski, University Clinic for Eye Diseases, Skopje, Republic of North Macedonia. E-mail: dr.isjanovski@gmail.com

Received: 2-Mar-2022; **Revised:** 20-Apr-2022;

Accepted: 27-Apr-2022; **Published:** 23-Jun-2022

Copyright: © 2022, Igor Isjanovski, Bisera Velkovska. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Competing Interests: The author have declared that no competing interests

ПРИКАЗ НА СЛУЧАЈ

СИНДРОМ НА “MORNING GLORY”-

Игор Исјановски¹, Бисера Велковска¹¹ Клиника за очни болести; Универзитет Св. Кирил и Методиј во Скопје, Медицински факултет, Република Северна Македонија**Извадок**

Синдромот на “Morning Glory” (MGS -Morning Glory Syndrom) е невообичаена конгенитална аномалија на оптичкиот нерв кој наликува на цвет познат како „morning glory“, го нарушува видот и може да биде поврзан и со окуларни и неокуларни абнормалности. Има карактеристичен фундоскопски изглед кој се состои од голема празнина во форма на инка на оптичкиот диск. Во овој труд презентираме невообичаена вродена аномалија на оптичкиот диск кај три годишно дете од женскиот пол.

Цитирање: Исјановски И, Велковска Б. Синдром на “Morning Glory”- приказ на случај Арх Ј Здравје 2022;14(1) 157-161.

doi.org/10.3889/aph.2022.6048

Клучни зборови: Синдром на “Morning Glory”, оптички диск.

***Кореспонденција:** Игор Исјановски, Универзитетска клиника за очни болести, Скопје, Република Северна Македонија. E-mail: dr.isjanovski@gmail.com

Примено: 2-мар-2022; **Ревидирано:** 20-апр-2022;

Прифатено: 27-апр-2022; **Објавено:** 23-јун-2022

Печатарски права: ©2022 Игор Исјановски, Бисера Велковска. Оваа статија е со отворен пристап дистрибуирана под условите на нелицензирана лиценца, која овозможува неограничена употреба, дистрибуција и репродукција на било кој медиум, доколку се цитираат оригиналниот(ите) автор(и) и изворот.

Конкурентски интереси: Авторот изјавува дека нема конкурентски интереси.

Introduction

Morning Glory Syndrome (MGS), first described by Kindler¹, is a congenital defect, a malformation of the optic nerve that resembles a flower known as morning glory. It is characterized by an enlarged funnel-shaped cavity in the optic disc, the part of the eye where the optic nerve fibers leave the retina. The disc is enlarged and has a white center (which gives the appearance of a white pupil). The number of blood vessels that are twisted is increased because they come from the enlarged disc that has a white center, which results in this malformation to look like the petals of a flower². In the medical literature, most reported and documented cases are unilateral (affecting only one eye), sporadic (without other cases in the family), and occur in females³⁻⁶.

Symptoms include very poor eyesight with poor visual acuity. MGS can occur on its own or in combination with other eye abnormalities, such as strabismus or lazy eye (amblyopia) or other non-ocular problems such as brain disorders⁷⁻⁹. Typically, individuals with non-ocular findings also have a broad head, a suppressed nasal bridge, and a defect or cleft in the middle of the upper lip⁹. MGS appears to be caused by the failure of the optic nerve to fully form as the baby develops. Manschot considers MGS to be a mesoderm disorder¹⁰. According to Lee & Traboulsi MGS is not inherited and the genetic defects associated with it have not been confirmed¹¹.

The most severe complication is retinal detachment and can occur in about 26-38% of people with MGS¹². MGS is sometimes misdiagnosed as optic nerve coloboma⁸.

The prevalence of MGS is unknown, a study conducted in Stockholm, Sweden registered a prevalence of 22.6 / 100,000⁵.

Treatment involves surgery and may result in some vision recovery. Depending on other related abnormalities, some patients require referral to several specialties such as neurosurgery, interventional neuroradiology, otolaryngology, and dentistry. All people diagnosed with "morning glory" disc anomaly should have a CT scan and MRI of the head³⁻⁴.

Case report

Child 3 years old, girl, was scheduled at the Department of Pediatric Ophthalmology at the Clinic for Eye Diseases Skopje with Dg.OD Esotropia cum hypertropia by a specialist ophthalmologist from secondary health care.

During taking the anamnesis from the girl's mother, it is found out that the curvature of the right eye inside the nose is from birth. Pregnancy was normal and the baby was born on time.

During the examination, the child is visibly upset and it is not possible to determine the visual acuity of both eyes and to examine the fundus, but the curvature of the right eye is visible.

Due to the age of the child, it was decided to examine the fundus under general anesthesia in the operating room with the help of a fundus camera.

An examination of the fundus under anesthesia diagnoses Morning Glory Syndrome of the right eye.

Bottom of the right eye: Papilla nervi optici (optic disc) enlarged to pale white. There are no blood vessels

through the gliosis tissue. Macula Lutea (point of clear vision) has a clear reflex. The retina is neat (Picture 1).

Left eye's left eye: Papilla nervi optici is at the level of the retina with clear boundaries, blood vessels with normal lumen and fullness. Macula Lutea has a clear reflex

Therapy: try to close the better eye (left eye) for 3 hours consecutively per day.

Advice is given for ENT examination and consultation with a neurophthalmologist.



Picture 1. Fundus of the right eye

Discussion

The etiology of the Morning Glory anomaly is poorly understood. There are some similarities with coloboma on the optical disc.

The visual prognosis in individuals with an anomaly of morning glory is poor. In addition to the abnormal disc itself and the propensity for serous retinal detachment, there is an added variable of complexity of high refractive errors, amblyopia, and strabis-

mus. Although surgery for eye strabismus and treatment of anisometric amblyopia is recommended and may result in some vision restoration, dramatic improvements are rare⁵. Treatment of other related abnormalities requires an interdisciplinary approach, often including neurosurgery, interventional neuroradiology, otolaryngology, and dentistry. All patients who are found to have a disc anomaly on a routine ophthalmologic examination should have a MRI of

the brain, MRI, and timely referrals to an appropriate subspecialist.

Conclusion

Patients with MGS need to be explained the need to wear goggles for two compelling reasons. Because the individual with MGS usually has beneficial vision in only one eye, preserving normal eye vision is essential. The strong link between MGS and retinal detachment also means that contact sports should be avoided, and if this is not possible, goggles designed to reduce the risk of ocular trauma should be worn. Educating the patient about the symptoms of retinal detachment will also be helpful as emergency medical care can prevent the detachment from worsening and allow for careful monitoring of the condition.

References

1. Kindler P. Morning glory syndrome: unusual congenital optic disc anomaly. *Am J Ophthalmol.* 1970;69(3):376-384.
2. Cohen MI, Matos CS, Amorim RLO, Ferreira WS, Cohen JM, Cohen MJ, Cohen SVA, Martins HO, Couto ISA, Souza MRS, Santos ALS. - Morning Glory Syndrome. Case Report and Literature Review, *J Brasileiro de Neurocirurg* 27 (4): 301 - 306, 2016
3. Ceynowa DJ, Wickström R, Olsson M, Ek U, Eriksson U, Wiberg MK et al. Morning Glory Disc Anomaly in childhood—a population-based study. *Acta ophthalmol.* 2015; 93:626-634. 3.
4. Fei P, Zhang Q, Li J, Zhao P. Clinical characteristics and treatment of 22 eyes of morning glory syndrome associated with persistent hyperplastic primary vitreous. *Br J Ophthalmol.* 2013; 97:1262-1267.
5. Itakura T, Miyamoto K, Uematsu Y, Hayashi S, Komai N. Bilateral morning glory syndrome associated with sphenoid encephalocele: case report. *J neurosurg.* 1992; 77:949-951.
6. Osaguona VB, Momoh RO., Morning Glory Syndrome in a Nigerian - A Case Report, *Journal of the West African College of Surgeons* 2017; 7(1):128-34.
7. Muslubas IS, Hocaoglu M, Arf S & Karacorlu M. A case of morning glory syndrome associated with persistent hyperplastic primary vitreous and Peters' anomaly. *GMS Ophthalmology Cases.* 2017; 7:Doc02.
8. Morning Glory Syndrome. Orphanet. 2017. Available at:http://www.orpha.net/consor/cgi-bin/Disease_Search.php?lng=EN&data_id=10399.
9. Strominger MB. Morning Glory Syndrome. *American Academy of Ophthalmology.* 2015; <https://www.aaof.org/pediatric-center-detail/neuro-ophthalmology-morning-glory-syndrome>.
10. Manschot WA. Morning Glory Syndrome: a histopathological study. *Br J Ophthalmol.* 1990;74(1):56-58.
11. Lee BJ, Traboulsi EI. Update on the morning glory disc anomaly. *Ophthalmic genet.* 2008; 29:47-52.
12. Lytvynchuk LM, Glittenberg CG, Ansari-Shahrezaei S & Binder S. Intraoperative optical coherence

tomography assisted analysis of pars Plana vitrectomy for retinal detachment in morning glory syndrome: a case report. *BMC Ophthalmology* 2017;17:134.